

Chiral Lagrangian from gauge invariant, nonlocal, dynamical quark model

Hua Yang^{a,b}, Qing Wang^{a,c}, Qin Lu^a

^a*Department of Physics, Tsinghua University, Beijing 100084, China **

^b*Institute of Electronic Technology, Information Engineering University, Zhengzhou 450004*

^c*Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China*

Abstract

Parameters of Gasser Leutwyler chiral Lagrangian are proved saturated by dynamical quark self energy $\Sigma(k^2)$ in a gauge invariant, nonlocal, dynamical quark model.

Much of low-energy QCD can be encoded into a series parameters appearing in a chiral Lagrangian, expanded to some finite order of low energy expansion. Attempts have been made to understand these parameters: it is shown that low lying vector mesons will saturate the parameters [1]. To go beyond phenomenological level, the anomaly contribution was taken as the main source of the parameters [2], we call this type investigation the anomaly approach, it leads result

$$8L_1 = 4L_2 = -2L_3 = 24L_7 = -8L_8 = L_9 = -2L_{10} = \frac{N_c}{48\pi^2} \quad (1)$$

which are close to experiment result except L_7 and L_8 which have wrong signs. The deficiency of this calculation lies in its independence of interaction: if we switch off the strong interaction and discuss a system of free quark field with external sources, the anomaly calculation can still be performed without any change. Then it seems that (1) is not due to strong interaction among quarks and gluons, but rather an artificial result. Another type research, we call it dynamical approach, mainly consider the dynamical effect [3], in which the main source of the parameters is from dynamical quark self energy $\Sigma(k^2)$. This approach has advantage of maintaining chiral symmetry and momentum dependence of dynamical quark

*Mailing address

mass, in the mean time avoiding introduce in the theory the hard constituent quark mass to cause wrong bad ultraviolet behavior of the theory. But it does not explain why it can offer the better numerical result (without wrong sign problem for L_7 and L_8) than anomaly approach. In fact, anomaly contribution and dynamical quark self energy contribution are two independent sources, if the anomaly contribution play role, according to (1), it will be dominant at all parameters and then there is no room left for $\Sigma(k^2)$ to play role to match the experiment data, except for L_7 and L_8 . If the anomaly contribution donot play role, it must be cancelled in some sense and after the cancellation, we need to show the remaining dynamical effect (which may or may not be dominant by dynamical quark self energy) can still recover or improve the result (1). It is purpose of this work to judge these two possibilities. We will show the second choice is correct, the cancellation do happen in dynamical approach and remanent contribution from dynamical quark self energy $\Sigma(k^2)$ can provide values for parameters of chiral Lagrangian consistent with experiment data.

Consider QCD in presence of external scalar,pseudoscalar, vector and axial vector sources,

$$J(x) = \not{p}(x) + \not{A}(x)\gamma_5 - s(x) + ip(x)\gamma_5,$$

The generating functional in Minkovski space is

$$Z[J] = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \mathcal{D}\Psi \mathcal{D}\bar{\Psi} \mathcal{D}A_\mu e^{i \int d^4x [\mathcal{L}(\psi, \bar{\psi}, \Psi, \bar{\Psi}, A_\mu) + \bar{\psi} J \psi]} \quad (2)$$

where ψ, Ψ, A_μ are light, heavy and gluon fields respectively. $\mathcal{L}(\psi, \bar{\psi}, \Psi, \bar{\Psi}, A_\mu)$ is Lagrangian of QCD. The chiral Lagrangian relate this generating functional by

$$Z[J] = \int \mathcal{D}U e^{i S_{\text{GL}}[U, J]} \quad (3)$$

U is pseudo goldstone boson (PGB) field, $S_{\text{GL}}[U, J]$ is Gasser and Leutwyler (GL) chiral Lagrangian [4],

$$S_{\text{GL}}[U, J] = S_{\text{normal}}[U, J] + S_{\text{anomaly}}[U, J], \quad (4)$$

$$\begin{aligned} S_{\text{normal}}[U, J] = \int d^4x \Big\{ & \frac{1}{4} F_0^2 \text{tr}[\nabla^\mu U^\dagger \nabla_\mu U + U \chi^\dagger + U^\dagger \chi] + L_1 [\text{tr}(\nabla^\mu U^\dagger \nabla_\mu U)]^2 \\ & + L_2 \text{tr}[\nabla_\mu U^\dagger \nabla_\nu U] \text{tr}[\nabla^\mu U^\dagger \nabla^\nu U] + L_3 \text{tr}[(\nabla^\mu U^\dagger \nabla_\mu U)^2] \\ & + L_4 \text{tr}[\nabla^\mu U^\dagger \nabla_\mu U] \text{tr}[\chi^\dagger U + U^\dagger \chi] + L_5 \text{tr}[\nabla^\mu U^\dagger \nabla_\mu U (\chi^\dagger U + U^\dagger \chi)] \\ & + L_6 [\text{tr}(\chi^\dagger U + U^\dagger \chi)]^2 + L_7 [\text{tr}(\chi^\dagger U - U^\dagger \chi)]^2 + L_8 \text{tr}[\chi^\dagger U \chi^\dagger U + \chi U^\dagger \chi U^\dagger] \\ & - i L_9 \text{tr}[F_{\mu\nu}^R \nabla^\mu U \nabla^\nu U^\dagger + F_{\mu\nu}^L \nabla^\mu U^\dagger \nabla^\nu U] + L_{10} \text{tr}[U^\dagger F_{\mu\nu}^R U F^{L, \mu\nu}] \\ & + H_1 \text{tr}[F_{\mu\nu}^R F^{R, \mu\nu} + F_{\mu\nu}^L F^{L, \mu\nu}] + H_2 \text{tr}[\chi^\dagger \chi] \Big\} + O(p^6) \text{ terms} \end{aligned} \quad (5)$$

$$S_{\text{anomaly}}[U, J] = S_{\text{WZW}}[U, J] + O(p^6) \text{ terms}, \quad (6)$$

where

$$\chi(x) = 2B_0[s(x) + ip(x)] \quad (7)$$

and $S_{\text{WZW}}[U, J]$ is Wess-Zumino-Witten action given in Ref. [5]. Up to order of p^4 , $S_{\text{anomaly}}[U, J]$ is completely known, but $S_{\text{normal}}[U, J]$ left fourteen parameters $F_0, B_0, L_1, \dots, L_{10}, H_1, H_2$ need to be calculated. To reveal the source of these parameters, we improve the conventional dynamical approach by building up a gauge invariant, nonlocal, dynamical (GND) quark model. The action in GND model is assumed to be $S_{\text{eff}}[\psi, \bar{\psi}, U, J]$, it relate to our generating functional by

$$Z[J] = \int \mathcal{D}U \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{iS_{\text{eff}}[\psi, \bar{\psi}, U, J]} . \quad (8)$$

The r.h.s. of above equation can be seen as a result of integrating out heavy quark and gluon fields and integrate in the PGB field U in (2). If we further integrate out light quark field in above generating functional, we obtain GL result (3). So $S_{\text{eff}}[\psi, \bar{\psi}, U, J]$ can be seen as an intermediate stage action to relate fundamental QCD with phenomenological chiral Lagrangian.

Compare (3) and (8), we find GL chiral Lagrangian relate to GND model by

$$e^{iS_{\text{GL}}[U, J]} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{iS_{\text{eff}}[\psi, \bar{\psi}, U, J]} , \quad (9)$$

$S_{\text{eff}}[\psi, \bar{\psi}, U, J]$ is required to be invariant under following local $U_L(3) \otimes U_R(3)$ chiral transformations:

$$\begin{aligned} \psi(x) &\rightarrow \psi'(x) = [V_R(x)P_R + V_L(x)P_L]\psi(x) \\ J(x) &\rightarrow J'(x) = [V_R(x)P_L + V_L(x)P_R][J(x) + i\cancel{D}][V_R^\dagger(x)P_R + V_L^\dagger(x)P_L] \\ U(x) &\rightarrow U'(x) = V_R(x)U(x)V_L^\dagger(x) . \end{aligned} \quad (10)$$

Notice that U field has standard decomposition $U(x) = \Omega(x)\Omega(x)$ and $\Omega(x)$ field, under transformation (10), transform as $\Omega(x) \rightarrow \Omega'(x) = h^\dagger(x)\Omega(x)V_L^\dagger(x) = V_R(x)\Omega(x)h(x)$ with $h(x)$ depend on V_R, V_L and Ω , represent an induced hidden local $U(3)$ symmetry to keep transformed Ω be a representative element at coset class.

To implement local chiral symmetry explicitly, we take a special local chiral transformation $V_R(x) = \Omega^\dagger(x), V_L(x) = \Omega(x)$, the corresponding hidden symmetry transformation is $h(x) = 1$,

$$\psi_\Omega(x) = [\Omega^\dagger(x)P_R + \Omega(x)P_L]\psi(x) \quad (11)$$

$$\begin{aligned} J_\Omega(x) &= [\Omega(x)P_R + \Omega^\dagger(x)P_L] [J(x) + i\cancel{D}] [\Omega(x)P_R + \Omega^\dagger(x)P_L] \\ &\equiv -s_\Omega(x) + ip_\Omega(x)\gamma_5 + \cancel{D}\Omega(x) + \cancel{D}\Omega(x)\gamma_5 \end{aligned} \quad (12)$$

$$U_\Omega(x) = 1 .$$

On rotated basis, we can rewrite (9) as

$$\begin{aligned}
e^{iS_{GL}[U,J]} &= \frac{\int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{iS_{\text{eff}}[\psi_\Omega, \bar{\psi}_\Omega, 1, J_\Omega]}}{\int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{i \int d^4x \bar{\psi}(x)[i\partial_x + J(x)]\psi(x)}} \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{i \int d^4x \bar{\psi}(x)[i\partial_x + J(x)]\psi(x)} \\
&= N' \frac{\int \mathcal{D}\psi_\Omega \mathcal{D}\bar{\psi}_\Omega e^{iS_{\text{eff}}[\psi_\Omega, \bar{\psi}_\Omega, 1, J_\Omega]}}{\int \mathcal{D}\psi_\Omega \mathcal{D}\bar{\psi}_\Omega e^{i \int d^4x \bar{\psi}_\Omega(x)[i\partial_x + J_\Omega(x)]\psi_\Omega(x)}}
\end{aligned} \tag{13}$$

where $N' \equiv \int \mathcal{D}\psi \mathcal{D}\bar{\psi} e^{i \int d^4x \bar{\psi}(x)[i\partial_x + J(x)]\psi(x)} = \text{Det}[i\partial_x + J(x)]$. In the last equality, we have taken chiral rotation (11) for functional integration measure both in numerator and denominator. The possible anomalies caused by this rotation are cancelled between numerator and denominator. Since we are only interested in U dependence of the theory, pure source terms N' is irrelevant and therefore can be treated as a normalization factor.

Result (13) tells us that S_{eff} should has following structure

$$S_{\text{eff}}[\psi_\Omega, \bar{\psi}_\Omega, 1, J_\Omega] = \int d^4x \bar{\psi}_\Omega(x)[i\partial_x + J_\Omega(x)]\psi_\Omega(x) + S_{\text{int}}[\psi_\Omega, \bar{\psi}_\Omega, 1, J_\Omega] \tag{14}$$

where S_{int} is interaction part caused by color gauge interaction. If we switch off color gauge interaction which means we are dealing with free fermion fields, there will be no effective Lagrangian ($S_{\text{GL}} = 0$). S_{int} should include those fermion self interaction terms caused by integrate out gluon and heavy quark fields in underlying QCD and integrate in local goldstone boson fields U . Among these, the most important effect related to chiral symmetry at low energy region is spontaneous chiral symmetry breaking (SCSB) which require quark has a nontrivial momentum dependent self energy $\Sigma(k^2)$, its effects can be introduced into the theory by adding in S_{int} a self energy term

$$- \int d^4x \bar{\psi}_\Omega(x) \Sigma(\partial_x^2) \psi_\Omega(x) . \tag{15}$$

Just this term itself is not enough, since it is not invariant under local chiral symmetry transformations. To make it invariant, in conventional dynamical approach [3], a non-integratable face factor is introduced into theory which cause very complex formulae and authors in [3] even donot put their analytical result in their papers. We donot use non-integratable face factor, instead we note that local chiral symmetry transformation on rotated variable is

$$\begin{aligned}
\psi_\Omega(x) &\rightarrow \psi'_\Omega(x) = h^\dagger(x) \psi_\Omega(x) \\
J_\Omega(x) &\rightarrow J'_\Omega(x) = h^\dagger(x) [J_\Omega(x) + i\partial_x] h(x) .
\end{aligned} \tag{16}$$

Original local chiral symmetry now is realized as a hidden local symmetry. Once the theory is constructed to be invariant under this hidden symmetry, it is invariant under original local chiral symmetry. Since interaction part in S_{eff} should be invariant on local chiral

symmetry, we need at least to generalize self energy term (15) to be invariant on hidden local symmetry (16). To achieve this, we change the ordinary derivative ∂_x^μ to hidden symmetry covariant derivative $\bar{\nabla}_x^\mu = \partial_x^\mu - iv_\Omega^\mu(x)$ (the overline on ∇_x^μ is to denote the difference with covariant derivative appeared in (5)). (16) tells us $v_\Omega(x)$ transform as $v_\Omega^\mu(x) \rightarrow v_\Omega^{\mu'}(x) = h^\dagger(x)v_\Omega^\mu(x)h(x) + ih^\dagger(x)[\partial^\mu h(x)]$ which lead $\bar{\nabla}_x^\mu \rightarrow \bar{\nabla}_x^{\mu'} = h^\dagger(x)\bar{\nabla}_x^\mu h(x)$. The modified chiral invariant interaction action now is

$$S_{\text{int}}[\psi_\Omega, \bar{\psi}_\Omega, 1, J_\Omega] = - \int d^4x \bar{\psi}_\Omega(x) \Sigma(\bar{\nabla}_x^2) \psi_\Omega(x). \quad (17)$$

This action is not the complete part of interaction, but it is the minimal part of interaction which respect local chiral symmetry with dynamical quark and SCSB. If we take the idea of dynamical perturbation originally from Pagel-Stokar [6] and developed in Ref. [7], in which at the leading order of the expansion, all perturbative effects are ignored and only nonperturbative effect considered in the theory is that from quark self energy $\Sigma(k^2)$. (17) in this sense can be seen as a result of leading order expansion from dynamical perturbation.

In GND model, quark fields dependence is bilinear and can be exactly integrated out, the result GL Lagrangian from (13) is

$$S_{GL}[U, J] \approx S_{GND}[U, J] \equiv -i\text{Tr} \ln[i\partial + J_\Omega - \Sigma(\bar{\nabla}^2)] + i\text{Tr} \ln[i\partial + J_\Omega]. \quad (18)$$

Use the Schwinger proper time formulation developed in [8], we can compute the $\Sigma(\bar{\nabla}^2)$ dependent determinant in (18). The result is

$$\begin{aligned} & -i\text{Tr} \ln[i\partial + J_\Omega - \Sigma(\bar{\nabla}^2)] \\ &= \int d^4x \text{tr}_f \left[B_0 F_0^2 s_\Omega + \mathcal{C}_1 a_\Omega^2 + \mathcal{C}_2 [d_\mu a_\Omega^\mu]^2 + \mathcal{C}_3 (d^\mu a_\Omega^\nu - d^\nu a_\Omega^\mu) (d_\mu a_{\Omega,\nu} - d_\nu a_{\Omega,\mu}) \right. \\ & \quad + \mathcal{C}_4 [a_\Omega^2]^2 + \mathcal{C}_5 a_\Omega^\mu a_\Omega^\nu a_{\Omega,\mu} a_{\Omega,\nu} + \mathcal{C}_6 s_\Omega^2 + \mathcal{C}_7 p_\Omega^2 + \mathcal{C}_8 s_\Omega a_\Omega^2 + \mathcal{C}_9 V_\Omega^{\mu\nu} V_{\Omega,\mu\nu} + \mathcal{C}_{10} V_\Omega^{\mu\nu} a_{\Omega,\mu} a_{\Omega,\nu} \\ & \quad \left. + \mathcal{C}_{11} p_\Omega d_\mu a_\Omega^\mu \right] + O(p^6) + \text{imaginary terms} \end{aligned} \quad (19)$$

where tr_f is trace for flavor indices. Covariant derivative for function f and $V_\Omega^{\mu\nu}$ are defined as

$$d^\mu f \equiv \partial^\mu f - iv_\Omega^\mu f + i f v_\Omega^\mu \quad V_\Omega^{\mu\nu} = \partial^\mu v_\Omega^\nu - \partial^\nu v_\Omega^\mu - iv_\Omega^\mu v_\Omega^\nu + iv_\Omega^\nu v_\Omega^\mu, \quad (20)$$

The Σ dependence for coefficients appeared in (19) are

$$F_0^2 B_0 = 4 \int d\tilde{k} \Sigma_k X_k \quad (21)$$

$$\mathcal{C}_1 = 2 \int d\tilde{k} \left[(-2\Sigma_k^2 - k^2 \Sigma_k \Sigma'_k) X_k^2 + (2\Sigma_k^2 + k^2 \Sigma_k \Sigma'_k) \frac{X_k}{\Lambda^2} \right] \quad (22)$$

$$\begin{aligned}
\mathcal{C}_2 &= -2 \int d\tilde{k} \left[-2A_k X_k^3 + 2A_k \frac{X_k^2}{\Lambda^2} - A_k \frac{X_k}{\Lambda^4} + \frac{k^2}{2} \Sigma_k^{\prime 2} \frac{X_k}{\Lambda^2} - \frac{k^2}{2} \Sigma_k^{\prime 2} X_k^2 \right] \\
\mathcal{C}_3 &= - \int d\tilde{k} \left[-2B_k X_k^3 + 2B_k \frac{X_k^2}{\Lambda^2} - B_k \frac{X_k}{\Lambda^4} + \frac{k^2}{2} \Sigma_k^{\prime 2} \frac{X_k}{\Lambda^2} - \frac{k^2}{2} \Sigma_k^{\prime 2} X_k^2 \right] \\
\mathcal{C}_4 &= 2 \int d\tilde{k} \left[\left(\frac{4\Sigma_k^4}{3} - \frac{2k^2 \Sigma_k^2}{3} + \frac{k^4}{18} \right) (6X_k^4 - \frac{6X_k^3}{\Lambda^2} + \frac{3X_k^2}{\Lambda^4} - \frac{X_k}{\Lambda^6}) + (-4\Sigma_k^2 + \frac{k^2}{2}) (-2X_k^3 \right. \\
&\quad \left. + \frac{2X_k^2}{\Lambda^2} - \frac{X_k}{\Lambda^4}) - \frac{X_k}{\Lambda^2} + X_k^2 \right] \\
\mathcal{C}_5 &= \int d\tilde{k} \left[\left(\frac{-4\Sigma_k^4}{3} + \frac{2k^2 \Sigma_k^2}{3} + \frac{k^4}{18} \right) (6X_k^4 - \frac{6X_k^3}{\Lambda^2} + \frac{3X_k^2}{\Lambda^4} - \frac{X_k}{\Lambda^6}) + 4\Sigma_k^2 (-2X_k^3 + \frac{2X_k^2}{\Lambda^2} \right. \\
&\quad \left. - \frac{X_k}{\Lambda^4}) + \frac{X_k}{\Lambda^2} - X_k^2 \right] \\
\mathcal{C}_6 &= 2 \int d\tilde{k} \left[(3\Sigma_k^2 + 2k^2 \Sigma_k \Sigma_k') X_k^2 + [-2\Sigma_k^2 - k^2(1 + 2\Sigma_k \Sigma_k')] \frac{X_k}{\Lambda^2} \right] \\
\mathcal{C}_7 &= 2 \int d\tilde{k} \left[(\Sigma_k^2 + 2k^2 \Sigma_k \Sigma_k') X_k^2 - k^2(1 + 2\Sigma_k \Sigma_k') \frac{X_k}{\Lambda^2} \right] \\
\mathcal{C}_8 &= 4 \int d\tilde{k} \left[(-4\Sigma_k^3 + k^2 \Sigma_k) X_k^3 + (4\Sigma_k^3 - k^2 \Sigma_k) \frac{X_k^2}{\Lambda^2} - (2\Sigma_k^3 - \frac{1}{2} k^2 \Sigma_k) \frac{X_k}{\Lambda^4} + 3\Sigma_k \frac{X_k}{\Lambda^2} - 3\Sigma_k X_k^2 \right] \\
\mathcal{C}_9 &= - \int d\tilde{k} \left[\left(\frac{1}{3} k^2 \Sigma_k' \Sigma_k'' + \frac{1}{3} \Sigma_k \Sigma_k'' \right) X_k + (C_k - D_k) \frac{X_k}{\Lambda^2} - (C_k - D_k) X_k^2 - 2E_k X_k^3 \right. \\
&\quad \left. + 2E_k \frac{X_k^2}{\Lambda^2} - E_k \frac{X_k}{\Lambda^4} \right] \\
i\mathcal{C}_{10} &= 4 \int d\tilde{k} \left[-2F_k X_k^3 + 2F_k \frac{X_k^2}{\Lambda^2} - F_k \frac{X_k}{\Lambda^4} + \frac{k^2}{2} \Sigma_k^{\prime 2} \frac{X_k}{\Lambda^2} - \frac{k^2}{2} \Sigma_k^{\prime 2} X_k^2 \right] \\
\mathcal{C}_{11} &= -4 \int d\tilde{k} \left[-(\Sigma_k + \frac{1}{2} k^2 \Sigma_k') \frac{X_k}{\Lambda^2} + (\Sigma_k + \frac{1}{2} k^2 \Sigma_k') X_k^2 \right] \tag{23}
\end{aligned}$$

where

$$\begin{aligned}
\int d\tilde{k} &\equiv iN_c \int \frac{d^4 k}{(2\pi)^4} e^{\frac{k^2 - \Sigma^2(-k^2)}{\Lambda^2}} & \Sigma_k &\equiv \Sigma(-k^2) & X_k &\equiv \frac{1}{k^2 - \Sigma^2(-k^2)} \\
A_k &= \frac{2}{3} k^2 \Sigma_k \Sigma_k' (1 + 2\Sigma_k \Sigma_k') + \frac{1}{3} \Sigma_k^2 (1 + 2\Sigma_k \Sigma_k') - \frac{1}{3} k^2 \Sigma_k^2 (\Sigma_k'^2 + \Sigma_k \Sigma_k'') + \frac{1}{6} k^4 (\Sigma_k'^2 + \Sigma_k \Sigma_k'') \\
B_k &= \frac{2}{3} k^2 \Sigma_k \Sigma_k' (1 + 2\Sigma_k \Sigma_k') + \frac{1}{3} \Sigma_k^2 (1 + 2\Sigma_k \Sigma_k') - \frac{1}{3} k^2 \Sigma_k^2 (\Sigma_k'^2 + \Sigma_k \Sigma_k'') + \frac{1}{18} k^4 (\Sigma_k'^2 + \Sigma_k \Sigma_k'') \\
&\quad + \frac{1}{6} k^2 (1 + 2\Sigma_k \Sigma_k') \\
C_k &= \frac{1}{3} - \frac{1}{3} \Sigma_k \Sigma_k' - \frac{1}{2} k^2 \Sigma_k'^2 \\
D_k &= \frac{1}{2} k^2 \Sigma_k'^2 + \frac{1}{6} k^2 \Sigma_k \Sigma_k'' (1 + 2\Sigma_k \Sigma_k') + \frac{2}{9} k^4 \Sigma_k' \Sigma_k'' (1 + 2\Sigma_k \Sigma_k') + \frac{2}{9} k^4 \Sigma_k'^2 (\Sigma_k'^2 + \Sigma_k \Sigma_k'') \\
&\quad + \frac{1}{3} k^2 \Sigma_k \Sigma_k' (\Sigma_k'^2 + \Sigma_k \Sigma_k'') \\
E_k &= -\frac{1}{6} k^2 \Sigma_k \Sigma_k' (1 + 2\Sigma_k \Sigma_k')^2 - \frac{1}{9} k^4 \Sigma_k'^2 (1 + 2\Sigma_k \Sigma_k')^2
\end{aligned}$$

$$F_k = -\frac{4}{3}k^2\Sigma_k\Sigma'_k + \frac{4}{3}k^2(\Sigma_k\Sigma'_k)^2 - \frac{2}{3}\Sigma_k^2 + \frac{2}{3}\Sigma_k^3\Sigma'_k - \frac{1}{3}k^2\Sigma_k^2(\Sigma_k'^2 + \Sigma_k\Sigma_k'') + \frac{1}{9}k^4(\Sigma_k'^2 + \Sigma_k\Sigma_k'') \\ + \frac{1}{3}k^2(1 + 2\Sigma_k\Sigma'_k) - \frac{1}{2}k^2.$$

The result for $\mathcal{C}_1 \equiv F_0^2$ in (22) is just the well known Pagel-Stokar formula [6], if we take momentum cutoff Λ be infinity. The part of $\Sigma(\overline{\nabla}^2)$ independent quark determinant in (19) is just the result of anomaly approach with a total minus sign, we can get result of anomaly approach by taking limit of $\Sigma_k = \text{const} \rightarrow 0$ (Note due to possible infrared divergence, limit of $\Sigma \rightarrow 0$ must be taken after the momentum integration). The nonzero coefficients $\mathcal{C}_i$ for the case of infinite momentum cutoff Λ is

$$\mathcal{C}_2 \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{24\pi^2}, \quad \mathcal{C}_3 \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{48\pi^2}(\ln \frac{\Sigma^2}{\Lambda^2} + \gamma + 1), \quad \mathcal{C}_4 \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{24\pi^2}(\ln \frac{\Sigma^2}{\Lambda^2} + \gamma + 4), \\ \mathcal{C}_5 \xrightarrow{\Sigma \rightarrow 0} -\frac{N_c}{24\pi^2}(\ln \frac{\Sigma^2}{\Lambda^2} + \gamma + 2), \quad \mathcal{C}_6 \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{8\pi^2}\Lambda^2, \quad \mathcal{C}_7 \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{8\pi^2}\Lambda^2 \\ \mathcal{C}_9 \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{48\pi^2}(\ln \frac{\Sigma^2}{\Lambda^2} + \gamma), \quad i\mathcal{C}_{10} \xrightarrow{\Sigma \rightarrow 0} \frac{N_c}{12\pi^2}(\ln \frac{\Sigma^2}{\Lambda^2} + \gamma + 2).$$

The pure imaginary terms in (19) are completely known at phenomenological level and its calculation in terms of Σ is already performed in Ref. [9] and proved exactly recover the Witten's result [5], we donot explicitly write down their detail structures.

With help of (12), (18) and (19) lead relation,

$$L_1 = \frac{1}{2}L_2 = \frac{\overline{\mathcal{C}}_5}{32} - \frac{\overline{\mathcal{C}}_9}{16} + i\frac{\overline{\mathcal{C}}_{10}}{32}, \quad L_3 = \frac{\overline{\mathcal{C}}_4 - 2\overline{\mathcal{C}}_5 + 6\overline{\mathcal{C}}_9 - 3i\overline{\mathcal{C}}_{10}}{16}, \\ L_4 = 0, \quad L_5 = \frac{\overline{\mathcal{C}}_8}{16B_0}, \quad L_6 = 0, \quad L_7 = \frac{\overline{\mathcal{C}}_2}{48} - \frac{\overline{\mathcal{C}}_{11}}{48B_0}, \\ L_8 = -\frac{\overline{\mathcal{C}}_2}{16} + \frac{\overline{\mathcal{C}}_6}{16B_0^2} - \frac{\overline{\mathcal{C}}_7}{16B_0^2} + \frac{\overline{\mathcal{C}}_{11}}{16B_0}, \quad L_9 = \frac{-4\overline{\mathcal{C}}_9 + i\overline{\mathcal{C}}_{10}}{8}, \\ L_{10} = \frac{-\overline{\mathcal{C}}_3 + \overline{\mathcal{C}}_9}{2}, \quad H_1 = \frac{\overline{\mathcal{C}}_3 + \overline{\mathcal{C}}_9}{4}, \quad H_2 = \frac{\overline{\mathcal{C}}_2}{8} + \frac{\overline{\mathcal{C}}_6}{8B_0^2} + \frac{\overline{\mathcal{C}}_7}{8B_0^2} - \frac{\overline{\mathcal{C}}_{11}}{8B_0}. \quad (24)$$

where $\overline{\mathcal{C}}_i \equiv \mathcal{C}_i - \lim_{\Sigma \rightarrow 0} \mathcal{C}_i$ $i = 1, 2, \dots, 11$. Term $-\lim_{\Sigma \rightarrow 0} \mathcal{C}_i$ is of special interest, since it relate to anomaly result mentioned in the beginning of this paper. In fact, in anomaly approach, the effective action is $i\text{Tr} \ln[i\hat{p} + J_\Omega]$ [2], which is just the result of (5) with (24) by taking $\mathcal{C}_i$ values at $\Sigma = 0$ and revert all signs. One can easily check this reproduce result (1) in which all ultraviolet divergence are cancelled each other for L_i parameters. The interpretation of this result is that from (18), *the contribution of anomaly play no role in the final result, it is completely cancelled by dynamical quark self energy dependent part, only the remainder after cancellation play role in the final parameters L_i .*

In (24), parameter B_0 needs special treatment. Since with help of (21), we find $F_0^2 B_0$ is generally divergent. To renormalize this condensate, we note that F_0 and $m_\Lambda \langle \overline{\psi}\psi \rangle_\Lambda$ (m_Λ is

bare current quark mass) is renormalization invariant or more general, the χ field defined in (7) is renormalization invariant, i.e. $\chi(x) = 2B_0[s(x) + ip(x)] = 2B_r[s_r(x) + ip_r(x)]$ with B_r, s_r, p_r are renormalized B_0, s, p . Correspondingly, renormalized quark condensate $\langle \bar{\psi}\psi \rangle_r$ is $\langle \bar{\psi}\psi \rangle_r = -N_f F_0^2 B_r$. So replacing the scalar and pseudoscalar sources and B_0 with renormalized ones donot change value of χ field. With renormalized sources, we can replace B_0 in (24) with B_r . In this paper, we donot directly calculate and use B_0 , instead we calculate and use B_r . The renormalization point is chosen to be at scale of 1GeV.

Now, once the quark self energy $\Sigma(k^2)$ was input into the formulae, we can get all parameters in Gasser-Leutwyler chiral Lagrangian. In conventional dynamical approach [3], the ignorance of $\Sigma(k^2)$ is parametrized by following ansatz

$$\Sigma(k^2) = \frac{(A+1)m^3}{k^2 + Am^2} \quad (25)$$

which satisfy $\Sigma(m^2) = m$ and shares qualitative similarities with solutions of improved ladder SD equation. It is finite, positive, monotonically decreasing functions with $1/k^2$ behavior at large q^2 and $\Sigma'(0) < 0$. The constituent quark mass m is determined for each choice of A for $F_0 = 93MeV$ from (22). For $A = 1, 2, 3, 4$, we obtain $m = 379, 350, 331, 317MeV$ respectively. With ansatz (25), the result parameters are listed in TABLE.I. We see that the wrong sign problem for L_7 and L_8 in anomaly calculation is corrected now and result parameters are roughly consistent with experiment data.

To further trace the relation of GND model with underlying theory QCD. Note that the parameters in the chiral Lagrangian are recently expressed in terms of QCD Green's functions [10] and for quark two point Green's function $\Phi_\Omega^{\sigma\rho}(x, y)$, at large N_c limit, [10] gives equation,

$$\begin{aligned} & [i\partial\!\!\!/ + i\Phi_\Omega^{T,-1} + J_\Omega + \tilde{\Xi}]^{\sigma\rho}(x, y) + \sum_{n=1}^{\infty} \int d^4x_1 \cdots d^4x_n d^4x'_1 \cdots d^4x'_n \frac{(-i)^{n+1}(N_c g^2)^n}{n!} \\ & \times \bar{G}_{\rho\rho_1 \cdots \rho_n}^{\sigma\sigma_1 \cdots \sigma_n}(x, y, x_1, x'_1, \cdots, x_n, x'_n) \Phi_\Omega^{\sigma_1\rho_1}(x_1, x'_1) \cdots \Phi_\Omega^{\sigma_n\rho_n}(x_n, x'_n) = 0. \end{aligned} \quad (26)$$

Now consider the coincidence limit of two point quark Green's function in QCD in presence of external sources,

$$\begin{aligned} & \frac{1}{N_c} \langle 0 | T \bar{\psi}_\alpha^{a\eta}(x) \psi_\alpha^{b\xi}(x') | 0 \rangle_{\text{QCD}} \equiv \frac{-i}{N_c} \frac{\delta \ln Z[J]}{\delta J^{(a\eta)(b\xi)}(x)} \\ & = \frac{1}{N_c} \frac{\int \mathcal{D}U \frac{\delta S_{\text{GL}}[U, J]}{\delta J^{(a\eta)(b\xi)}(x)} e^{iS_{\text{GL}}[U, J]}}{\int \mathcal{D}U e^{iS_{\text{GL}}[U, J]}} = \frac{\int \mathcal{D}U \Phi^{(a\eta)(b\xi)}[U, J](x, x) e^{iS_{\text{GL}}[U, J]}}{\int \mathcal{D}U e^{iS_{\text{GL}}[U, J]}} \end{aligned} \quad (27)$$

where we have used (3) and $\Phi^{(a\eta)(b\xi)}[U, J](x, x)$ is

$$\Phi^{(an)(b\xi)}[U, J](x, x) \equiv \frac{1}{N_c} \frac{\delta S_{\text{GL}}[U, J]}{\delta J^{(an)(b\xi)}(x)} \approx \frac{1}{N_c} \frac{\delta S_{\text{GND}}[U, J]}{\delta J^{(an)(b\xi)}(x)}. \quad (28)$$

Use (18), the rotated Φ become

$$\Phi_{\Omega}^{(an)(b\xi)}[U, J](x, x) = \text{Tr} \left[\frac{-i}{i\partial + J_{\Omega} - \Sigma(\bar{\nabla}^2)} \frac{\delta[J_{\Omega} - \Sigma(\bar{\nabla}^2)]}{\delta J_{\Omega}^{(an)(b\xi)}(x)} \right] \quad (29)$$

which imply

$$\Phi_{\Omega}(x, y) = -i \int d^4z [i\partial + J_{\Omega} - \Sigma(\bar{\nabla}^2)]^{-1}(x, z) [\delta(z - y) + \Delta(z, y)] \quad (30)$$

where $\Delta(z, y)$ relate to $\delta\Sigma(\bar{\nabla}^2)/\delta J$. Compare to (26), we find, present choice of S_{int} is equivalent to take following approximation

$$\begin{aligned} & \left[(1 + \Delta)^{-1} [\Sigma(\bar{\nabla}^2) - \Delta(i\partial + J_{\Omega})] \right]^{\sigma\rho}(x, y) \\ & \approx \tilde{\Xi}^{\sigma\rho}(x, y) + \sum_{n=1}^{\infty} \int d^4x_1 \cdots d^4x_n d^4x'_1 \cdots d^4x'_n \frac{(-i)^{n+1} (N_c g^2)^n}{n!} \\ & \quad \times \bar{G}_{\rho\rho_1 \cdots \rho_n}^{\sigma\sigma_1 \cdots \sigma_n}(x, y, x_1, x'_1, \cdots, x_n, x'_n) \Phi_{\Omega}^{\sigma_1\rho_1}(x_1, x'_1) \cdots \Phi_{\Omega}^{\sigma_n\rho_n}(x_n, x'_n). \end{aligned} \quad (31)$$

If we further drop correlation functions $\bar{G}_{\rho\rho_1 \cdots \rho_n}^{\sigma\sigma_1 \cdots \sigma_n}(x, y, x_1, x'_1, \cdots, x_n, x'_n)$ with $n > 1$ and ignore the external sources in above equation (Δ therefore must be ignored), as mentioned in Ref. [10], (31) then is just Schwinger-Dyson equation for quark propagator. We only consider the term of quark self energy with argument of $\bar{\nabla}^2$ which is the minimal generalization from pure $\Sigma(\partial^2)$ term to source dependent terms satisfying local chiral symmetry. Include in source terms, just self energy term in l.h.s. of (31) is not enough to match all contributions of r.h.s. of equation, but if underlying QCD can provide correct predictions for parameters in chiral Lagrangian, the fact of $\Sigma(k^2)$ dominance in the parameters of chiral Lagrangian imply that the terms we dropped at this work should not play so important role. we will leave these terms in addition to self energy term to balance equation (31) in future investigations.

In conclusion, all 12 parameters in p^4 order $SU(3)$ chiral Lagrangian are explicitly expressed in terms of functions of quark self energy $\Sigma(k^2)$. We have shown that the original result given from anomaly approach is completely cancelled. Take suitable quark self energy ansatz to perform numerical calculation, we find after cancellation of anomaly contribution, the dynamical quark self energy do can provide parameter values roughly consistent with experimental data.

ACKNOWLEDGMENTS

This work was supported by National Science Foundation of China No.90103008 and fundamental research grant of Tsinghua University.

REFERENCES

- [1] G.Ecker, J.Gasser,A.Pich and E.de Rafael, Nucl. Phys. **B321** (1989)311; J.F.Donoghue, C.Ramirez and G.Valencia, Phys. Rev. **D39** (1989)1947
- [2] J. Balog. Phys. Lett. B **149**, 197 (1984); A.A. Andrianov and L. Bonora, Nucl. Phys. B **233**, 232 (1984); A.A. Andrianov, Phys. Lett. B **157**, 425 (1985); A.A. Andrianov et al, Phys. Lett. B **186**, 401 (1987); D. Espriu, E. De Rafael, and J. Taron, Nucl. Phys. B **345**, 22 (1990).
- [3] B. Holdom, Phys. Rev. D **45**, 2534 (1992); B. Holdom, J. Terning, and K. Verbeek, Phys. Lett. B **245**, 612 (1990); B **273**, 549(E) (1991); B. Holdom, *ibid.* B **294**, 293 (1992)
- [4] J. Gasser and H. Leutwyler, Ann. Phys. **158**, 142 (1984); Nucl. Phys. B **250**, 465 (1985).
- [5] E.Witten, Nucl. Phys. **B223**, 422(1983).
- [6] H.Pagels and S.Stokar, Phys. Rev. **D20**, 2947(1979)
- [7] B. Holdom and Q.Wang, Phys. Rev. D **55**, 3101 (1997).
- [8] Qin Lu, Hua Yang and QingWang, hep-ph/0202078, to appear in Commun. Theor. Phys.
- [9] C.D.Roberts, R.T.Cahill and J.Praschifka, Ann. Phys. **188**, 20(1988).
- [10] Qing Wang, Yu-ping Kuang, Ming Xiao and Xuelel Wang, Phys. Rev. **D61**, 54011(2000);

TABLES

TABLE I. Values, multiplied by 10^3 , for the parameters of the order p^4 chiral Lagrangian calculated in GND model with quark self energy determined by ansatz (25). anomaly: anomaly approach result; expt: experimental values.

	L_1	L_2	L_3	L_4	L_5	L_6	L_7	L_8	L_9	L_{10}
A=1	0.927	1.85	-6.55	0	1.79	0	-0.570	1.56	3.79	-5.18
A=2	0.771	1.54	-5.53	0	1.63	0	-0.501	1.39	2.60	-3.91
A=3	0.708	1.42	-5.09	0	1.51	0	-0.449	1.26	2.13	-3.38
A=4	0.674	1.35	-4.85	0	1.41	0	-0.413	1.17	1.90	-3.08
anomaly	0.792	1.58	-3.17	0	0	0	0.263	-0.792	6.33	-3.17
expt	0.9 ± 0.3	1.7 ± 0.7	-4.4 ± 2.5	0 ± 0.5	2.2 ± 0.5	0 ± 0.3	-0.4 ± 0.15	1.1 ± 0.3	7.4 ± 0.7	-6.0 ± 0.7